

***Rhexoacrodictys broussonetiae* sp. nov. from Guizhou, China**

ZHONG-JIU XIAO^{1,2}, XIAO-XIA LI^{1,2*},
HAO-DONG WANG³, PEI-YONG SONG^{1,2}, LING TANG¹

¹College of Biological & Agricultural Science and Technology, Zunyi Normal College,
Zunyi, Guizhou 563002, China

²Key Laboratory of Regional Characteristics for Conservation &
Utilization of Plant Resource in Chishui River Basin, Zunyi, Guizhou, 563002, China

³Nursing Department Heyuan City Health School, Guangdong, 517000, China

*CORRESPONDENCE TO: xzj198099@163.com

ABSTRACT—A new hyphomycete, *Rhexoacrodictys broussonetiae* is described and illustrated from specimens collected on dead *Broussonetia papyrifera* stems in Guizhou Province, China. It is characterized by distinct, unbranched conidiophores with monoblastic, percurrently extending conidiogenous cells that produce acrogenous, subspherical to spherical, 17–28 × 19–31 µm, 0–2 transversely septate conidia with several additional oblique longitudinal septa. A dichotomous key to the four *Rhexoacrodictys* species is provided.

KEY WORDS—*Acrodictys*, anamorphic fungi, *Ascomycota*, taxonomy

Introduction

Baker & Morgan-Jones (2002) established *Rhexoacrodictys* to accommodate four *Acrodictys* species: *R. erecta* (Ellis & Everh.) W.A. Baker & Morgan-Jones, *R. fimicola* (M.B. Ellis & Gunnell) W.A. Baker & Morgan-Jones, *R. fuliginosa* (B. Sutton) W.A. Baker & Morgan-Jones, and *R. queenslandica* (Matsush.) W.A. Baker & Morgan-Jones. Delgado (2009) later proposed the combination *R. martinii* (J.L. Crane & Dumont) G. Delgado, but Xia & al. (2017) transferred *R. martinii* to *Distoseptispora* and *R. queenslandica* to *Junewangia*, based on morphology and phylogenetic analysis. Thus, *Rhexoacrodictys* currently contains only three accepted species, which are mainly characterized by

indeterminate and regeneratively percurrently extending conidiogenous cells that often acquire a narrow circumscissile dehiscence zone at the extreme apex. The conidia, which are ellipsoid, ovoid, subspherical to spherical, transversely, longitudinally and obliquely septate, undergo rhexolytic conidial secession.

During ongoing surveys of saprobic microfungi from the subtropical forests of Guizhou Province, China, a *Rhexoacroditys* specimen collected on dead stems of *Broussonetia papyrifera* was found to differ from all accepted *Rhexoacroditys* taxa and is proposed here as a new species, *R. broussonetiae*. The holotype is deposited in the Mycological Herbarium of Zunyi Normal College, Zunyi, China (HMZNC).

***Rhexoacroditys broussonetiae* Xiao X. Li & Z.J. Xiao, sp. nov.**

FIG. 1

FUNGALNAME FN570519

Differs from all other *Rhexoacroditys* species by its subspherical to spherical conidia.

TYPE: China, Guizhou Province: Xishui National Nature Reserve, on dead stems of *Broussonetia papyrifera* (L.) Vent. (*Moraceae*), 27 Jun. 2017, X.X. Li (Holotype, HMZNC 0246).

ETYMOLOGY: in reference to the host genus, *Broussonetia*.

COLONIES on natural substrate effuse, hairy, brown to dark brown. Mycelium partly superficial, partly immersed in the substratum, composed of branched, septate, pale brown to brown, cylindrical, smooth-walled hyphae, 2–4 µm diam. CONIDIOPHORES macronematous, mononematous, single or a loose cluster of 2–4, erect, straight or slightly flexuous, cylindrical, smooth, thick-walled, dark brown to blackish, 4–7-septate, 70–119 × 4.5–6.5 µm, inflated toward the base, 6.5–9 µm diam, blackish, tapering slightly, paler brown, indeterminate. CONIDIOGENOUS CELLS integrated, terminal, monoblastic, with a pale brown, narrow circumscissile dehiscence zone at the extreme apex, 3–4.5 µm diam, percurrently extending. Conidial secession rhexolytic. CONIDIA holoblastic, solitary, acrogenous, subspherical to spherical, with 0–2 transverse septa and several longitudinal and oblique septa, with darkly-pigmented bands over the septa, smooth, brown to dark brown, 17–28 × 19–31 µm, bearing a small marginal frill at the base, 2.5–4.5 µm diam, pale brown.

COMMENTS—*Rhexoacroditys broussonetiae* produces conidia that are similar in shape to those in *Distoseptispora martinii* (J.L. Crane & Dumont) J.W. Xia & X.G. Zhang, which differ in their (sub)globose to broadly ellipsoidal shape and smaller size (11–16 µm diam.—Delgado 2009; 15–20 × 11–16 µm—Xia & al. 2017). *Rhexoacroditys broussonetiae* also differs from the other three *Rhexoacroditys* species, which produce ellipsoid or ovoid conidia.




FIG. 1. *Rhexoacrodictys broussonetiae* (holotype, HMZNC 0246)
A. Conidiophores with conidia; B. Conidiophores and conidiogenous cells; C. Conidia.

Key to *Rhexoacrodictys* species

- 1. Conidia dictyosporous, subglobose to globose
(17–28 × 19–31 µm, 0–2 transverse septa) *R. broussonetiae*
- 1. Conidia dictyosporous, ellipsoid or ovoid 2
- 2. Conidia without a dark brown apex, more or less evenly pigmented
(15–24 × 11–15 µm, 2–3 transverse septa, medium to blackish
brown) *R. fimicola*
- 2. Conidia apically dark brown to black, darker upwards when mature 3
- 3. Conidia 24–40 × 15–22 µm,
septa predominantly oblique, dark brown or black *R. erecta*
- 3. Conidia 21–26 × 12–16 µm,
septa 3(–4) transverse, brown to blackish brown *R. fuliginosa*

Acknowledgments

The authors express gratitude to Dr. Xiu-Guo Zhang and Dr. Jian Ma for serving as pre-submission reviewers and for their valuable comments and suggestions. Dr. Lorelei L. Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated. This project was supported by the National Natural Science Foundation of China (Nos. 31600030) and the Science and Technology Project of Guizhou Province (No. LKZS2012-18, LKZS2012-25, LKZS2014-20).

Literature cited

Baker WA, Partridge EC, Morgan-Jones G. 2002. Notes on hyphomycetes LXXXVII. *Rhexoacrodictys*, a new segregate genus to accommodate four species previously classified in *Acrodictys*. Mycotaxon 82: 95–113.

Delgado G. 2009. South Florida microfungi: *Veramycella bispora*, a new palmicolous, anamorphic genus and species, with some new records for the continental USA. Mycotaxon 107: 357–373. <https://doi.org/10.5248/107.357>

Xia JW, Ma YR, Li Z, Zhang XG. 2017. *Acrodictys*-like wood decay fungi from southern China, with two new families *Acrodictyceae* and *Junewangiaceae*. Scientific Reports 7:7888 [21 p.]. <https://doi.org/10.1038/s41598-017-08318-x>